


1/11

FIGURE 1




DEST AVAILABLE COPY







4/11



5/11



6/11



7/11

FIGURE 7

FBBC Consultant: Foodstuff and Botanicals Benefit and Consumption Consultant

	Detailed inform		
	Detailed information		
	Common Name 15 Tomato		
	Specific name: Amico tomato		
	Synonym Pseudomonas syringae toma		
	Area of origin Godollo, Hungary		
	State of sample Raw and fresh ()		
阿拉拉克	Benifits and Consum	THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER OF THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER.	
	Health benefits or inedical effect	Recommended Minimum \\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	
	Anti-oxidant, Anti-aging	Fo 101 og/day of Fig. 3, 5 Fig. 1	
	Ann-cancer (preventing prostate, lung, and stomach cancers)	169.g/day	
	2 2 2 2 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3		
	Processed or cooked tomato provides more!	Senerius (
		60-	1
	<i></i>	· 603	
	501		

8/11

FIGURE 8

707

709

701

9/11


FIGURE 9

FBICD:

Foodstuff and Botanical Ingredient and Content Database

Detailed Information (Bl@Dide alled lillormation Common Name Ginkgo Latin Name Ginkgo biloba L Indigeñous Yin xin', Ginkgo, Gingko Name Nan Xiong, Guang Dong Province, China Collection Site Oct 1994 Collection Time State Powder. Part for Analysis Leaf. To aid circulatory problems in the elderly, especially insufficiency and the consequent cognitive effects, peripheral circulatory impairment, particularly intermittent claudication (poor circulation to the lower legs), and vertigo and Health or. Medical Effects tinnitus. To protect against altitude sickness and to mediate erectile dysfunction in males. To treat cerebral insufficiency such as Alzheimer's disease. Analysis Method HPLC Compound Flavonoids Class **Total Class** 13.2mg/g Content Chi,Jing Rui, et al. Acta Pharmaceutica Sinica, 1997, 32(8) 625-628 ingrédiénis and Content Compound Name Content (of leaf) quercetin; 0.15 mg/g kaempferol 1.3 mg/g bilobetin 1.0 mg/g ginkgetin 4.2 mg/g sciadoptisin 6.5 mg/g

10/11



11/11

FIGURE 11

EDC database: Effective Dose of Chemicals

	Detailed Information
	Detailed Informations of the second second
Chemical Nam	e Cucurbitacin B
Synonym	Amarın, Fabaceın II
CAS No Saurce	6199-67-3 Towel gourd, melon
Reference	Ji, YB & Zhang GM, Pharmacological Action and Application of Available Antitumon Composition of Traditional Chinese Medicine, HelongJiang Science and Technology publishing
	Effective Dose
	Effect Minimum Effective dose on the second
Anti-cancer	lin vitro; ED50 = 0:005 μg/ml
Anti-cancer	in.wv::0:25-0.5mg/kg
Stomach wash	
Hypo-injecti	on (mouse) LD50 = (1,0±0:07)mg/kg



This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

© BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
☐ FADED TEXT OR DRAWING
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
COLOR OR BLACK AND WHITE PHOTOGRAPHS
☐ GRAY SCALE DOCUMENTS
☐ LINES OR MARKS ON ORIGINAL DOCUMENT
☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
□ other:

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.